

Targeting Tumor-Associated Macrophages by Anti-tumor Chinese Materia Medica*

PU Wei-ling¹, SUN Li-kang^{1,2,3}, GAO Xiu-mei^{1,2}, Curzio Rüegg⁴, Muriel Cuendet⁵, Micheal O. Hottiger⁶,
ZHOU Kun^{1,2}, MIAO Lin^{1,2,6}, ZHANG Yun-sha^{3,4}, and Margaret Gebauer⁷



Prof. SUN Li-kang

ABSTRACT Tumor-associated macrophages (TAMs) play a key role in all stages of tumorigenesis and tumor progression. TAMs secrete different kinds of cytokines, chemokines, and enzymes to affect the progression, metastasis, and resistance to therapy depending on their state of reprogramming. Therapeutic benefit in targeting TAMs suggests that macrophages are attractive targets for cancer treatment. Chinese materia medica (CMM) is an important approach for treating cancer in China and in the Asian region. According to the theory of Chinese medicine (CM) and its practice, some prescriptions of CM regulate the body's internal environment possibly including the remodeling the tumor microenvironment (TME). Here we briefly summarize the pivotal effects of TAMs in shaping the TME and promoting tumorigenesis, invasion, metastasis

and immunosuppression. Furthermore, we illustrate the effects and mechanisms of CMM targeting TAMs in anti-tumor therapy. Finally, we reveal the CMM's dual-regulatory and multi-targeting functions on regulating TAMs, and hopefully, provide the theoretical basis for CMM clinical practice related to cancer therapy.

KEYWORDS tumor-associated macrophages, tumor microenvironment, Chinese materia medica, Chinese medicine, anti-tumor therapy

Infiltration of the tumor microenvironment (TME) by immune and inflammatory cells is involved in promoting tumorigenesis, tumor progression, local invasion and metastasis.⁽¹⁾ Tumor-associated macrophages (TAMs), a prominent and abundant inflammatory component of solid and hematological malignancies, express a pro- and anti-tumor effect in all stages of carcinogenesis. The molecular and functional phenotype of TAMs plays a key role in the tumorigenesis and progression.⁽²⁾ A malignant tumor contains not merely proliferating neoplastic cells, but many other cells which constitute a microenvironment, for example, endothelial cells, fibroblasts, and infiltrating immune cells. TAMs, one of the important infiltrating immune cells, secrete an array of cytokines, chemokines, and enzymes that affect tumor progression, metastasis, and resistance to therapy.^(3,4) TAMs secrete an array of cytokines, chemokines, and enzymes that affect tumor progression, metastasis, and resistance to therapy.⁽⁵⁻⁷⁾ Clinical success in targeting macrophage (MΦ) in tumor patients suggests that TAMs might be attractive targets in combination therapy for tumor treatment.^(8,9)

complementary medicine parallel to the conventional medicine system for tumor patients in China and other Asian countries. Chinese materia medica (CMM) is the natural therapeutic agent used under the guidance of CM theories.⁽¹⁰⁾ CMM therapies are a long-term clinical practice, and share the characteristics of

Chinese medicine (CM) is an important

1. Key Laboratory of Pharmacology of Traditional Chinese Medicine Formulae, Ministry of Education, Tianjin University of Traditional Chinese Medicine, Tianjin (300193), China; 2. Tianjin State Key Laboratory of Modern Chinese Medicine, Tianjin University of Traditional Chinese Medicine, Tianjin (300193), China; 3. School of Integrative Medicine, Tianjin University of Traditional Chinese Medicine, Tianjin (300193), China; 4. Pathology Unit, Department of Medicine, Faculty of Sciences, University of Fribourg, Fribourg, Switzerland; 5. School of Pharmaceutical Sciences, University of Geneva, University of Lausanne, Genève, Switzerland; 6. Department of Molecular Mechanisms of Disease, University of Zurich-Irchel, Zurich, Switzerland; 7. Chingcura, Center for Traditional Chinese Medicine, Zurich, Switzerland
Correspondence to: Prof. SUN Li-kang, Tel: 86-22-59596163, Fax: 86-22-59596163, E-mail: likang.sun@bluewin.ch
DOI: <https://doi.org/10.1007/s11655-017-2974-y>

multi-component, multi-targeted and dual-regulatory function.⁽¹¹⁾ One of the mechanisms of CMM tumor therapy is related to regulating the internal environment and remodeling the TME, which coincides with conventional tumor therapy targeting the TME. CMM prescriptions, consisting of a single herb or combination of several herbs based on the theory of CM, have been used for treating and preventing various cancers during long-term popular practice or in combination therapy with chemotherapeutics.^(7,12,13) The potential mechanism of anti-tumor effects of CMM includes increasing the efficacy of the therapeutic effect of chemotherapeutics including decreasing the development of chemoresistance or regulating the immune system.⁽¹⁴⁻¹⁶⁾ It has been shown that CMMs have the effects of enhancing the immunological surveillance, regulating the body's internal milieu and remodeling the TME.⁽¹³⁾

In this article, we briefly introduce the origin, subpopulations, phenotypic characteristics and polarization of TAM and the TAM's effect on the TME in promoting tumorigenesis, metastasis, and immune suppression. Specifically we analyzed the effects and mechanisms of CMM prescriptions, extracts or active compounds on targeting TAMs in anti-tumor therapies. We also discuss the CMM's dual-regulatory and multi-targeting functions on regulating TAM, and hopefully, provide the theoretical basis for CMM use in clinical practice for cancer therapy.

Origin, Subpopulation and Polarization of TAM

MΦs involved in the immune responses appear to generate from circulating bone marrow monocytes, and are recruited to the TME by cytokines secreted by the tumor cells and other cells, then differentiated into various phenotypes and functions including anti- or pro-tumor activities.^(5,17,18)

Subpopulations of MΦ

In *in vitro* models, MΦs have been classified as pro-inflammatory, classically activated MΦs (M1) and anti-inflammatory; alternatively activated MΦs (M2) based on their cell surface phenotypes and secreted cytokines.

M1-type MΦs, induced by Th1 mediators such as interferon (IFN)- γ , tumor necrosis factor α (TNF- α) or lipopolysaccharides (LPS), release pro-inflammatory cytokines such as interleukin (IL)-12, IL-23. They express major histocompatibility complex (MHC) II molecules, and have a high IL-12/IL-10 secretion ratio.

M1 MΦs participate in the Th1 inflammatory responses primarily through production of pro-inflammatory mediators and up-regulation of cell surface molecules necessary for antigen presentation to activate Th1 cells and enhance the ability to phagocytose pathogenic material.^(19,20) M2-type MΦs, typically induced by Th2 cytokines, mediate resistance to parasites, mediate immune regulation and tissue repair.^(4,21) M2 MΦs release high levels of IL-10 and low levels of IL-12 and IL-23. In the context of cancer, however, M2 MΦs help driving tumor growth through the promotion of angiogenic programs, tissue remodeling, and directly or indirectly suppress cytotoxic T cell activity.⁽²²⁾

The classification of MΦs in type M1 versus type M2 is an oversimplification of the *in vivo* situation. Most recently, Mosser and colleagues⁽²³⁾ suggested a new grouping of MΦ populations based on fundamental MΦ functions. This form of classification, though it may better reflect the heterogeneity and plasticity of MΦs in the TME, is clearly more difficult to consider and apply to research. The classical classification in M1/M2 MΦs is still commonly used as a valid and accepted concept for the sake of clarity.⁽²⁾

Polarization of MΦ

MΦs can be phenotypically polarized into diverse functional phenotypes by different microenvironment, and could have different effects during early or late tumor progression.⁽²⁴⁾ Tumor escape has been linked to a polarization switch from M1 to M2 MΦ phenotypes. In the early steps of tumor initiation, TAMs were mostly polarized towards a M1-like phenotype while during later steps of tumor progression they polarized mostly toward a M2-like phenotype.^(24,25) In contrast, Wang, et al⁽²⁶⁾ reported a pro-metastatic role of M1-like TAM (CD68⁺ HLA-DR⁺) in hepatocellular carcinoma. This difference may be due to the different and complex environments of various cancers observed *in vivo*, which cannot yet be easily simulated *in vitro*. Many chemokines, such as C-X-C motif chemokine 12 (CXCL12),⁽²⁷⁾ and chemokine ligand 2 (CCL2),⁽²⁸⁾ signaling pathways, such as nuclear factor κ B (NF- κ B),⁽²⁹⁾ Notch receptors,⁽³⁰⁾ or hypoxia present in the TME⁽²⁸⁾ contribute to the polarization of TAM. The details of TAM function and CMM targeting MΦs will be presented below.

Effects of TAMs during Carcinogenesis and Anti-tumor Mechanisms of CMM Targeting TAMs

Many observations indicate that TAMs possess

M2-associated pro-tumor functions, including the induction of tumor angiogenesis and the stimulation of tumor cell proliferation, survival and invasion by producing inflammatory mediator, such as cytokines, and matrix metalloprotease (MMPs), or suppressing the cytotoxic effect of T cells.^(2,31) TAMs play an important role in growth, metastasis and immunosuppression. Recently, studies showed that "TAM-specific therapies" may be a suitable option for further therapeutic approaches in cancer. Thus, CMM targeting TAMs may be considered as an additional anti-tumor therapy applicable to multiple stages of carcinogenesis.^(19,32)

Tumor Initiation

It is now widely accepted that the nonmalignant cells of the TME evolve along with the growing tumor and provide essential cues supporting the modulation of TAM phenotype.⁽⁵⁾ Chronic inflammation was shown to be associated with the development of malignant tumors by creating a tumor-promoting microenvironment.⁽³³⁾ MΦs are recruited to different areas of the TME and educated toward different phenotypes.^(17,18) It has been shown that pancreatic cancer cells could activate MΦs *in vitro* to polarize toward an M2-phenotype, through the cytokines IL-4, IL-10 and CCL2.^(34,35) This effect was confirmed by *in vivo* results revealing about 90% of all TAMs to be M2 MΦs. Accordingly, depletion of MΦs *in vivo* resulted in a significantly reduction in tumor growth.⁽³⁶⁾ Some Chinese herbal ingredients or prescriptions have been confirmed to effectively inhibit the recruitment and activation of MΦs to tumor sites and to re-educate TAM (see below).

Recruitment and Activation of TAMs

Diverse cytokines and polypeptides produced by tumor cells as well as microenvironment, such as hypoxia, are involved in guiding the recruitment, positioning and function of TAMs in the TME.⁽³⁾ For example, CCL2, plays a pivotal role in many cancers, including pancreas, ovary, and breast in the recruitment and infiltration on MΦ into the TME.^(34,35,37) In a pancreatic cancer mouse model, the inhibition of CCL2 was shown to reduce the recruitment of TAMs into the tumor, to improve the response to gemcitabine and increase the antitumor T-cell response and further inhibit tumor metastasis.^(37,38)

CMM may regulate the recruitment and activation of MΦs in the TME by various pathways. Berberine, an isoquinoline alkaloid isolated from many medicinal herbs, such as *Coptis chinensis* Franch, inhibited the growth of CT26 tumor cells implanted subcutaneously

in BABL/c mice and significantly reduced the density of CD68⁺/CD206⁺ TAMs. This is consistent with the notion that the anti-tumor effect of berberine involves inhibition of TAMs recruitment.^(39,40) Epigallocatechin gallate (EGCG), a polyphenol present in green tea, inhibited tumor growth in a murine breast cancer model by inhibiting TAM infiltration and M2 MΦ polarization.^(41,42) Praeruptorin D and E, the main constituents of *Peucedanum praeruptorum* Dunn, could protected BALB/c mice from lipopolysaccharide (LPS)-induced acute lung injury by inhibiting MΦs infiltration to the lung, cytokine release and vascular permeability.⁽⁴³⁾ Polysaccharopeptide (PSP), from *Coriolus versicolor* demonstrated anti-tumor and immune regulating effects.^(44,45) PSP significantly increased the number of CD14⁺/CD16⁺ monocytes in phytohemagglutinin treated peripheral blood mononuclear cell model, but could not affect the proliferations of T-cells, natural killer, and B-cells. Thus, stimulating monocyte/MΦ with PSP could be an effective therapeutic intervention in targeting tumors.⁽⁴⁶⁾

Astragalus polysaccharides (APS), the polysaccharide fraction of the roots of *Astragalus membranaceus*, were reported to activate MΦs in a toll-like receptor 4 (TLR4)-dependent manner while activating B cells via membrane Ig in a TLR4-independent manner.⁽⁴⁵⁾ Abrus agglutinin is a low-toxicity protein from the plant *Abrus precatorius* L. In its native (NA) and heat-denatured (HDA) forms, it activated TAMs in Dalton's lymphoma-bearing mice, and significantly increased *in vitro* cytotoxicity against tumor cells and stimulated production of nitric oxide but not of TNF- α . Thus, NA and HDA could be used as regulators of TAMs, for the treatment of cancer.⁽⁴⁷⁾ KSG-002, a 30% ethanolic extract of *Astragalus membranaceus* and *Angelica Sinensis* gigas at 3:1 w/w ratio, suppressed xenograft tumor growth and pulmonary metastasis via inhibiting TAM recruitment by blocking NF- κ B-dependent TNF- α production.⁽⁴⁸⁾ Qinghao Biejia Decoction (青蒿鳖甲汤, QHBJ), a classical prescription consisting of *Artemisia apiacea*, turtle shell, *Rehmannia glutinosa* Libosch., *Anemarrhena asphodeloides* Bge. and *Paeonia suffruticosa* Andr, significantly reduced the incidence rate of dimethylbenzanthracene (DMBA)-induced ovarian cancer in rats, inhibited the infiltration of TAMs in ovarian tumors and down-regulated the expression of inflammatory mediators prostaglandin E₂ (PGE₂) and leukotriene B4 (LTB4) and the angiogenic

factor vascular endothelial growth factor (VEGF) in ovarian tumor tissues.⁽⁴⁹⁾ Shaoyao Decoction (芍药汤), a classical CMM prescription of *Paeonia lactiflora* Pall effective in treating ulcerative colitis, could significantly increase the survival of mice with azoxymethane (AOM)/dextran sodium sulfate (DSS)-induced colitis-associated colorectal cancer (CA-CRC) by reducing the infiltration of TAM and thereby decreasing the expression levels of serum IL-1 β , IL-6, and TNF- α .⁽⁴⁸⁾ PHY906 (KD018), a CMM prescription consisting of *Glycyrrhiza uralensis* Fisch, *Paeonia lactiflora* Pall, *Scutellaria baicalensis* Georgi, and *Ziziphus jujuba* Mill, may potentiate sorafenib action by up-regulating the expression of CCL2, which could attract M1-type M Φ s and exert an anti-tumor activity.⁽⁴⁹⁾

Re-programing of TAM Phenotype

Mantovani proposed that TAM exert pleiotropic functions that can influence tumor growth and progression in opposite directions.⁽³⁵⁾ Recently, researches confirmed that circulating monocytes were recruited to the TME by chemotactic factors produced by tumor cells and differentiated toward the M2 phenotype.⁽⁵²⁻⁵⁴⁾ M2 M Φ s release various growth factors and cytokines to promote tumor cell invasion, induce angiogenesis, suppress antitumor immunity, and accelerate tumor cell metastasis.^(55,56) The polarization of TAM was a regulated and reversible process and dependent on the TME. It was shown that a predominance activation of NF- κ B and signal transducers and activators of transcription (STAT) 1 promoted M1 M Φ polarization resulting in cytotoxic and inflammatory activities, while STAT3 and STAT6 activation resulted in M2 M Φ polarization leading to immune suppression and tumor progression.^(57,58)

CMM targeting TAMs could act as an anti-tumor prescription by aiming at preventing polarization of TAMs toward a more anti-tumor phenotype. Emodin, a natural anthraquinone derivative isolated from *Rheum palmatum* L., inhibited M Φ recruitment and M2 polarization by suppressing STAT6 phosphorylation and C/EBP β expression in the 4T1 and EO771 orthotopic breast cancer mouse models.^(59,60) Isoliquiritigenin (ISL), a flavonoid from *Glycyrrhiza uralensis* Fisch, inhibited M2 M Φ polarization by down-regulating PGE₂ and IL-6-mediated signaling in a mouse model of AOM/DSS-induced colitis-associated tumorigenesis. In parallel, ISL also blunted the M2 polarization of RAW264.7 cells and mouse

peritoneal M Φ s by inactivating PGE₂/peroxisome proliferator-activated receptor δ and IL-6/STAT3 signaling *in vitro*.⁽⁶¹⁾ Sinomenine, an alkaloid extracted from *Sinomenium Acutum*, has been used to treat rheumatism in China. Zhang, et al⁽⁶²⁾ found that treating breast cancer-bearing mice with 100 mg/kg sinomenine hydrochloride (SH) can provide a self-reinforcing stimulus for further polarizing TAMs away from an M2-like phenotype toward an M1-like phenotype thereby contributing to its anti-tumor and anti-metastasis effect. The aqueous extract of *Crinum latifolium* L. leaf induced expression of TNF- α , IL-1 β and IL-6 and promoted M Φ s differentiation into the pro-inflammatory M1 phenotype.⁽⁶³⁾ The *Sarcandra glabra* polysaccharide (SGP) could effectively increase expression of CD14, CD16/32 and CD40 in M Φ s induced by IFN- γ , which indicated the potential for applications in regulating the immune balance in the treatment of cancer.⁽⁶⁴⁾ Fuzheng Jiedu Decoction (扶正解毒汤, FZJD), a prescription consisting of *Astragalus membranaceus*, *Codonopsis pilosula*, *Atractylodes macrocephala*, *Lycium barbarum*, *Smilax glabra*, *Polygonum multiflorum*, and *Actinidia arguta*, reduced the expression of the M2 M Φ s specific genes Arg-1 (an immunosuppressive enzyme) and CCL22 (a chemotactic chemokine) in an *in vitro* co-culture system of RAW264.7 M Φ s and mesenchymal stem cells.⁽¹³⁾ FZJD combined with 5-fluorouracil (5-FU) inhibited the marker of M2 M Φ s subtype expression and increased the marker of M1 M Φ s subtype expression, resulting in the inhibition of tumor recurrence and metastasis in a gastric cancer model.⁽⁶⁵⁾

In some excessive inflammatory reactions, active CMM components could promote phenotype transition from M1 to M2, thus acting as anti-inflammation drug. Berberine down-regulated the content of TNF- α and the mRNA expression levels of inducible nitric oxide synthase (iNOS) and suppressor of cytokine signaling (SOCS) 3 in M Φ s induced by LPS, inhibited the content of IL-10 and the mRNA expression level of Arg-1 in M Φ s induced by IL-4, but did not modify SOCS2. The results indicate that berberine has an effect on inhibiting the M1 and M2 polarization of M Φ s *in vitro* and may play a regulatory role in the dynamic balance of M1/M2 phenotype.⁽⁶⁶⁾ The similar results were found in many compounds or CM prescription, such as curcumin,⁽⁶⁷⁾ arctigenin,⁽⁶⁸⁾ and Xuebijing Injection (血必净注射液).⁽⁵²⁾

Importantly, CMM can act in a dual-regulatory function in the inflammatory reaction depending on the

specific physiological or pathological processes and the tissues involved. CMM can suppress excessive inflammation, while promoting the polarization of anti-inflammatory M2 MΦs into M1 phenotype thereby altering the state of immuno-suppression in the TME of advanced cancers.

Tumor Metastasis

The process of metastasis represents a critical phase in the neoplastic cascade, and growing evidence indicate a tight link between metastasis and TAMs, whereby TAMs produce inflammatory cytokines, stimulate the growth of tumor cells and promote tumor cell migration and metastasis. TAMs contribute to tumor progression also by producing several factors enhancing angiogenesis and lymphogenesis.^(34,69)

TAMs have been reported to promote angiogenesis via the production of diverse pro-angiogenic factors such as transforming growth factor- β (TGF- β), VEGF, Platelet derived growth factor (PDGF), fibroblast growth factor (FGF) and angiogenic chemokines, such as CXCL12.^(70,71) They also regulate matrix degradation through the production of matrix-degrading enzymes such as MMPs, plasmin, and urokinase-type plasminogen activator (uPA).^(69,72) Clinical research has confirmed that M2-polarized TAMs are associated with poor prognoses in several cancers, including classical Hodgkin's lymphoma (CHL),⁽⁷³⁾ lung adenocarcinoma,⁽⁷⁴⁾ and pancreatic cancer.⁽⁷⁵⁾ Ding, et al⁽⁷⁶⁾ found a significant correlation between the expression of CD68 or CD163 protein with a tumor size, lymph node metastasis, and vascular invasion in patients with pancreatic cancer consistent with the role of TAM infiltration in pancreas and the promotion of pancreatic cancer development.

Angiogenesis and Lymphangiogenesis

Angiogenesis and lymphangiogenesis are critical processes for tumor growth, invasion, and metastasis. TAMs promote angiogenesis in a wide type of tumors, such as gliomas, carcinomas of the esophagus, breast, bladder, and prostate by releasing vascular growth factors (mainly VEGF and PDGF), cytokines (e.g. TGF- β), proteases, and chemokines.⁽⁷⁷⁾ TAMs support tumor lymphangiogenesis not only by secreting pro-lymphangiogenic factors, such as VEGF-A and MMP-9, but also by trans-differentiation into lymphatic endothelial cells.⁽⁷⁸⁾ Wogonin, a flavonoid isolated from the root

of *Scutellaria baicalensis*, inhibited tumor growth, metastasis, angiogenesis and lymphangiogenesis, and reduced the number of TAM in the tumors of LM8-bearing mice. Wogonin inhibited VEGF-C-induced lymphangiogenesis through the reduction of VEGF-C-induced VEGF receptor-3 phosphorylation.⁽⁷⁹⁾ Recently a phenolic alkaloids of *Menispermum dauricum* (PAMD) inhibited BXPC-3 tumor growth and angiogenesis by reducing the production of VEGF, and basic fibroblast growth factor (bFGF).⁽⁸⁰⁾ Many results show that genistein, a soy isoflavonoid, inhibited angiogenesis through the regulation of VEGF, MMPs, and epidermal growth factor receptor expressions, thereby resulting in a strong anti-tumor effect.^(81,82) Qingyi Huaji Decoction (清胰化积汤, QYHJ), consists of many herbs including the radics of *Paeonia lactiflora* Pall and *Scutellaria baicalensis* Georgi, has been used as an integrative agent in the clinical treatment of advanced pancreatic cancers, especially those with liver metastasis. QYHJ inhibited the growth of human pancreatic cancer cell line CFPAC-1 *in vivo*, and this correlated with the down-regulation of serum level of pro-angiogenic factors sVCAM-I, TNF- α , bFGF, IL-6 and IL-8.⁽⁸³⁾ Besides, QYHJ exerted an inhibitory effect on the growth of experimental liver metastasis in nude mice inoculated intra-splenically with the human pancreatic cancer cell line SW1990, in part though the inhibition of VEGF and Cyr61 expression.⁽⁸⁴⁾ QYHJ inhibited cancer-related inflammation in the SW1990HM (pancreatic tumor cell line with high metastasis) tumor model by decreasing infiltration of TAM and IL-6 production and resulting in the decreased epithelial-to-mesenchymal transition and cell invasion.⁽⁸⁵⁾ Hongjingtian Injection (红景天注射液), a CMM standardized Tibetan herbal product extracted from *Rhodiola crenulata* L., used to treat patients with acute ischemic stroke or coronary heart disease, inhibited the metastasis of human non-small cell lung cancer by reducing the production of VEGF.⁽⁸⁶⁾

Matrix Remodeling

In the tumor stroma, MΦs can produce enzymes promoting matrix degradation, such as MMPs, uPA and uPA receptor, and plasmin, thereby facilitating invasion and metastasis of tumor cell.⁽³⁴⁾ Emodin exerts anti-proliferative and anti-metastatic activity in pancreatic cancer both *in vitro* and *in vivo*, which may be related to the down-regulation of NF- κ B and its regulated molecules survivin and MMP-9.⁽⁸⁷⁾ Gallic acid and gallic acid dimmer, polyphenols extracted

from pericarp of water caltrop, significantly decreased MMP-9 activity and inhibited migration of gastric cancer cells; TNF- α and IL-6 associated with migration and angiogenesis were also significantly reduced.⁽⁸⁸⁾ Aloe emodin, an anthraquinone from *Rheum palmatum* L., has anti-migratory and anti-angiogenic activities in colon cancer cells by down-regulating expression of MMP-2/9, RhoB and VEGF.⁽⁸⁹⁾ Honokiol, a major phenolic compound isolated from the root and stem bark of magnolia, inhibited the expression of MMP-9 in PANC-1 cells *in vitro* and VEGF in tumor tissue *in vivo* in an athymic mouse tumor model.⁽⁹⁰⁾

Immune Suppression

TAMs also produce various immune suppressive factors, including PGE₂, indoleamine 2,3-dioxygenase, Arg-1, and IL-10 to suppress tumor-associated antigen-specific T cell immunity and thus contribute to immunosuppression.^(31,34) The major immune stimulatory cytokine IL-12 may be due to the activity of IL-10, produced either by TAM or by the tumor cells. Part of the immune suppressive activity of TAM is exerted indirectly by their release of chemokines, such as CCL18 and CCL22 that preferentially attract T cell subsets devoided of cytotoxic functions.⁽⁹¹⁾ These immune stimulatory cytokines and chemokines could be the potential targets of anti-tumor CMM. Different reports indicate that polysaccharides from CMM herbs could improve the amount of M Φ s in the TME, enhance phagocytosis, as well as stimulate the secretion of NO, TNF- α , IL-1, etc., possibly related with M1 M Φ s activation and polarization.⁽⁹²⁾

Direct Effect on M Φ s

PAC-I, a mannose-rich polysaccharide fraction from *Aloe vera* L. var. *chinensis* (Haw) Berg., increased the expression of MHC-II and Fc γ R, and enhanced endocytosis, phagocytosis, NO production, TNF- α secretion and tumor cell cytotoxicity *in vitro*, stimulated systemic TNF- α production *in vivo* and prolonged the survival of tumor-bearing mice. The results may have therapeutic implications of PAC-I in tumor immunotherapy.⁽⁹³⁾

Regulation of Immune System

Salidroside, an active component of *Rhodiola rosea* L., promoted proliferation and phagocytosis of peritoneal M Φ s stimulated by LPS and IFN γ , and reduced the production of reactive oxygen species, but promoted the production of NO, and in activated peritoneal

M Φ s.⁽⁹⁴⁾ Li, et al⁽⁹⁵⁾ demonstrated that berberine inhibited the proliferation of Ishikawa endometrial carcinoma cancer cells co-cultured with TAM and the production of IL-8, while blocking NF- κ B signaling pathway to restrain metastasis formation. Cucurbitacin D, an oxygenated tetracyclic triterpenoid isolated from *Trichosanthes kirilowii*, increased LPS-induced IL-1 β production in culture supernatant of M Φ s, indicating that cucurbitacin D could initiate immune regulating activity in M Φ s and possibly function as anti-tumor drug.⁽⁹⁶⁾ 20S-dihydroprotopanaxadiol, a protopanaxadiol derivative found in *Panax ginseng* exhibiting various biological activities including anti-cancer, pro-apoptotic, and anti-inflammatory effects, boosted innate immune responses of monocytes and M Φ s.⁽⁹⁷⁾ Garlic lectin was shown to induce IL-12 production via activation of p38 MAPK/ERK in mouse M Φ s and to stimulate IFN- γ production in spleen cells, consistent with anti-tumor effects.⁽⁹⁸⁾ Many studies showed that polysaccharides could enhanced cytotoxic and phagocytic activities of immune cells and increased the productions of many Th1 cytokines via various signaling pathways, such as *Panax ginseng* Polysaccharide,⁽⁹⁹⁾ *B. striata* Polysaccharide,^(100,101) *Polyporus* polysaccharide,^(102,103) *Plantago asiatica* L. Polysaccharide,⁽¹⁰⁴⁾ *Ganoderma lucidum* Polysaccharide.⁽¹⁰⁵⁾ These effects induced anti-cancer effect indirectly though the activation of immune cells rather than via direct cytotoxic effects.^(106,107) Water-soluble polysaccharide extracted from *Sanguisorba officinalis* L., significantly inhibited the growth of mouse transplanted tumor, remarkably increased the spleen index and promoted splenocytes proliferation, M Φ phagocytosis and secretion of serum cytokines IL-2, TNF- α and IFN- γ in sarcoma 180-bearing mice in the absence of direct cytotoxic activity on tumor cells. These results indicated that the anti-tumor activity of SOP may be achieved through immune potentiation.⁽¹⁰⁸⁾ These results were confirmed by other researchers using different CMM, such as polysaccharide from *Limonium sinense* Kuntze,⁽¹⁰⁹⁾ *Lilium* sp.,⁽¹¹⁰⁾ *G. lucidum*⁽¹¹¹⁾ and Weikangfu Granule (胃康复胶囊), a drug preparation used to treat gastric cancer and consisting of *Curcuma wenyujin*, *Astragalus membranaceus*, *Glycyrrhiza inflata*, *Poria coco*,⁽¹¹²⁾ and Shenqi Fuzheng Injection (参芪扶正注射液), a CMM standardized product extracted from radices *Codonopsis* sp. and *Astragali* sp. to improve immune function against chronic diseases.⁽¹¹³⁾ PSSC, polysaccharides from *Salvia chinensis* Benth, suppressed growth of H22 cells and reduced serum levels of PGE₂ in mice, suggesting an anti-tumor immune-stimulatory

activity.⁽¹¹⁴⁾ Two basidiomycete species, *Lentinus edodes* and *Cordyceps sinensis*, increased IL-2 production in LBRM-33 1A5 T cells, through indirect action on MΦs, without directly acting on T cell.⁽¹¹⁵⁾ Qingshu Yiqi Decoction (清暑益气汤), a CMM prescription consisting of nine herbs including *Astragalus membranaceus*, *Panax ginseng* and *Scutellaria baicalensis*, has been classically used for treating fever and pulmonary disorders. Tumor burden and losses of muscle mass were found to be significantly decreased when 5-FU was combined with this Chinese prescription, suggesting that the combination of *S. baicalensis* and QYD is able to ameliorate cachectic symptoms and positively stimulate anti-tumor immunity in mice undergoing chemotherapy.⁽¹¹⁶⁾

Conclusion

CM, respectively CMM, belongs to a natural medicine system, which is still in active practice as an alternative therapeutic possibility beyond the origin cultural region. Based on this medicine system, many new medications have been developed, such as compounds isolated from *Artemisia annua*, and minnelide (a potential anti-pancreatic cancer medication, modified from triptolide).⁽¹¹⁷⁻¹¹⁹⁾

In the last two decades, experimental and clinical evidence showed that most chronic diseases, including cancer, are caused by a dysregulated inflammatory-immune response, and TAMs are a key link between inflammation and cancer in regulating tumor growth, progression and response to anticancer drugs.⁽¹²⁰⁾ CMM demonstrated obvious anti-tumor effects based on its characteristics of multi-components and multi-targets. The anti-cancer properties of CMM targeting TAMs were shown to involve the following mechanisms: (1) inhibition of the production of cytokines or chemokines associated with TAM recruitment and activation; (2) promotion of MΦs phenotype conversion from M2 to M1; (3) reduction of the production of pro-tumoral cytokines secreted by TAM including angiogenic and lymphangiogenic factors;⁽⁷⁸⁾ (4) reshaping of the TME to inhibit the immunosuppressive responses.

Most findings of anti-tumor CMM targeting MΦs or immune regulation are related to the expression level of cytokines or chemokines in the TME. More investigations are required, however, to unravel the mechanism involved and identify active components of CMM prescriptions. CMM prescriptions are highly complex systems, and interactions between different components

commonly exist.⁽¹²¹⁾ For example in the case of extracts of radices *Astragalus membranaceus* and *Angelica sinensis* in different proportions (e.g. w/w ratio) opposite effects related to cancer pulmonary metastasis were observed and should be investigated.⁽⁴⁸⁾

In this article, we have analyzed the literature for studies focused on cancer therapy related to CM and CMM. It is not just limited to active single substances, classical prescriptions, but it also includes combinations with conventional therapies, and the putative targeting phase in cancer development as listed in Figure 1. Further investigations are required to better characterize the anti-tumor effects of CM and CMM. In order to develop novel anti-tumor therapies based on CMM it will be essential to confirm results obtained in *in vitro* studies addressing molecular mechanisms of action on TAMs by performing *in vivo* studies on tumor progress.⁽¹²²⁾




Figure 1. Anti-tumor CMMs Targeting TAMs Effects on Tumor Progression

Notes: Pre-malignant or malignant cells and microenvironments, such as inflammation or hypoxia, generated various cytokines, chemokines, and enzymes to recruit and activate macrophages. The TAMs, interacted with TME and other cells, re-programmed into diverse functional phenotypes to affect tumor initiation, metastasis, and immune suppression. Anti-tumor CMM compounds or prescriptions targeting TAMs could effect on different phases of tumor growth through different mechanism.

REFERENCE

1. Ostuni R, Kratochvill F, Murray PJ, Natoli G. Macrophages and cancer: from mechanisms to therapeutic implications. Trends Immunol 2015;36:229-239.
2. Mantovani A, Schioppa T, Porta C, Allavena P, Sica A. Role of tumor-associated macrophages in tumor progression and invasion. Cancer Metast Rev 2006;25:315-322.
3. Sica A, Saccani A, Mantovani A, Allavena P, Sica A. Tumor-associated macrophages: a molecular perspective. Int immunopharmacol 2002;2:1045-1054.
4. Schmieder A, Michel J, Schonhaar K, Goerdts S, Schledzewski K. Differentiation and gene expression profile of tumor-associated macrophages. Semin Cancer Biol 2012;22:289-297.
5. Noy R, Pollard JW. Tumor-associated macrophages: from mechanisms to therapy. Immunity 2014;41:49-61.

6. Cook J, Hagemann T. Tumour-associated macrophages and cancer. *Curr Opin Pharmacol* 2013;13:595-601.
7. Gao J, Tongchuan H, Yingbo L, Wang YT. A traditional Chinese medicine formulation consisting of *Rhizoma Corydalis* and *Rhizoma Curcumae* exerts synergistic anti-tumor activity. *Oncol Rep* 2009;22:1077-1083.
8. Shimura S, Yang G, Ebara S, Wheeler TM, Frolov A, Thompson TC. Reduced infiltration of tumor-associated macrophages in human prostate cancer: association with cancer progression. *Cancer Res* 2000;60:5857-5861.
9. Bolli E, Movahedi K, Laoui D, Van Ginderachter JA. Novel insights in the regulation and function of macrophages in the tumor microenvironment. *Curr Opin oncol* 2017;29:55-61.
10. Yang XW. Discovery strategy for effective and active constituents of Chinese material medica based on processes of metabolism and disposition in intra-body. *China J Chin Mater Med (Chin)* 2007;5:001.
11. Kyrn PS. On didirectionally regulated effects of traditional Chinese materia medica [dissertation]. Jinan: Shandong University of Traditional Chinese Medicine; 2003.
12. Cai Y, Luo Q, Sun M, Corke H. Antioxidant activity and phenolic compounds of 112 traditional Chinese medicinal plants associated with anticancer. *Life Sci* 2004;4:2157-2184.
13. Li J, Guo Q, Lin H. Regulating effects and molecular mechanisms of traditional Chinese medicine in the tumor immune suppressive environment. *World Chin Med (Chin)* 2014;9:845-850.
14. Zhao F, Liu P. Progress of study on action mechanisms of TCM in anti-tumor and preventing metastasis of tumor. *Chin J Integr Tradit West Med (Chin)* 2007;27:178-181.
15. Xu T, Xu RA. Progress on Anti-tumor mechanism of Chinese medicine and its active ingredients. *J Huaqiao Univ (Natural Sci, Chin)* 2009;30:360-365.
16. Zhang X, Zhou F. Anti-tumor traditional Chinese medicine research progress. *J Liaoning Univ Tradit Chin Med (Chin)* 2012;14:142-144.
17. Mantovani A, Allavena P, Sozzani S, Vecchi A, Locati M, Sica A. Chemokines in the recruitment and shaping of the leukocyte infiltrate of tumors. *Semin Cancer Biol* 2004;14:155-160.
18. Taddei ML, Giannoni E, Comito G, Chiarugi P. Microenvironment and tumor cell plasticity: an easy way out. *Cancer Letters* 2013;341:80-96.
19. Kennedy BC, Showers CR, Anderson DE, Anderson L, Canoll P, Bruce JN. Tumor-associated macrophages in glioma: friend or foe? *J Oncol* 2013;2013:486912.
20. Mueller CK, Schultze-Mosgau S. Histomorphometric analysis of the phenotypical differentiation of recruited macrophages following subcutaneous implantation of an allogeneous acellular dermal matrix. *Int J Oral Maxillof Surg* 2011;40:401-407.
21. Mantovani A, Allavena P. The interaction of anticancer therapies with tumor-associated macrophages. *J Exp Med* 2015;212:435-445.
22. Ruffell B, Affara NI, Coussens LM. Differential macrophage programming in the tumor microenvironment. *Trends Immunol* 2012;33:119-126.
23. Mosser DM, Edwards JP. Exploring the full spectrum of macrophage activation. *Nat Rev Immunol* 2008;8:958-969.
24. Martinez FO, Sica A, Mantovani A, Locati M. Macrophage activation and polarization. *Front Biosci* 2008;13:453-461.
25. Sun P, Wang Y. Tumor-associated macrophages development and treatment of tumor. *Cancer Res Prev Treat* 2014;41:829-833.
26. Wang H, Wang X, Li X, Fan YC, Li GS, Guo C, et al. CD68⁺HLA-DR⁺ M1-like macrophages promote motility of HCC cells via NF-kappaB/FAK pathway. *Cancer Letters* 2014;345:91-99.
27. Sanchez-Martin L, Estechea A, Samaniego R, Sanchez-Ramon S, Vega MA, Sanchez-Mateos P. The chemokine CXCL12 regulates monocyte-macrophage differentiation and RUNX3 expression. *Blood* 2011;117:88-97.
28. Rius J, Guma M, Schachtrup C, Akassoglou K, Zinkernagel AS, Nizet V, et al. NF-kappaB links innate immunity to the hypoxic response through transcriptional regulation of HIF-1alpha. *Nature* 2008;453:807-811.
29. Saccani A, Schioppa T, Porta C, Biswas SK, Nebuloni M, Vago L, et al. P50 nuclear factor-kappaB overexpression in tumor-associated macrophages inhibits M1 inflammatory responses and antitumor resistance. *Cancer Res* 2006;66:11432-11440.
30. Wang Y, He F, Feng F, Liu XW, Dong GY, Qin HY, et al. Notch signaling determines the M1 versus M2 polarization of macrophages in antitumor immune responses. *Cancer Res* 2010;70:4840-4849.
31. Jinushi M, Komohara Y. Tumor-associated macrophages as an emerging target against tumors: creating a new path from bench to bedside. *Bba-Rev Cancer* 2015;1855:123-130.
32. Galdiero MR, Bonavita E, Barajon I, Garlanda C, Mantovani A, Jaillon S. Tumor associated macrophages and neutrophils in cancer. *Immunobiology* 2013;218:1402-1410.
33. Pesic M, Greten FR. Inflammation and cancer: tissue regeneration gone awry. *Curr Opin Cell Biol* 2016;43:55-61.
34. Allavena P, Sica A, Solinas G, Porta C, Mantovani A. The inflammatory micro-environment in tumor progression: the role of tumor-associated macrophages. *Crit Rev Oncolhematol* 2008;66:1-9.
35. Mantovani A, Bottazzi B, Colotta F, Porta C, Mantovani A. The origin and function of tumor-associated macrophages. *Immunol Today* 1992;13:265-270.
36. Partecke LI, Gunther C, Hagemann S, Jacobi C, Merkel M, Sendler M, et al. Induction of M2-macrophages by tumour cells and tumour growth promotion by M2-macrophages: a quid pro quo in pancreatic cancer. *Pancreatol* 2013;13:508-516.
37. Mitchem JB, Brennan DJ, Knolhoff BL, Belt BA, Zhu Y, Sanford DE, et al. Targeting tumor-infiltrating macrophages decreases tumor-initiating cells, relieves immunosuppression, and improves chemotherapeutic responses. *Cancer Res* 2013;73:1128-1141.
38. Lunardi S, Muschel RJ, Brunner TB, Belt BA, Zhu Y, Sanford DE, et al. The stromal compartments in pancreatic cancer: are there any therapeutic targets. *Cancer Lett* 2014;343:147-155.
39. Li J, Hao Y, Zhang Y, Xie W, Li J, Xue XO. Berberine inhibits tumor-associated macrophages in subcutaneous tumor of mice. *Chin J Histochem Cytochem (Chin)* 2011;20:203-206.
40. Wang Y. Berberine exerts its anti-tumor effects by targeting cancer cells as well as interfering with the tumor microenvironment in prostate cancer [dissertation]. Jinan: School of Medicine, Shandong University; 2011.
41. Hong M, Wang N, Tan HY, Tsao SW, Feng YB. MicroRNAs and Chinese medicinal herbs: new possibilities in cancer therapy. *Cancers* 2015;7:1643-1657.
42. Jang JY, Lee JK, Jeon YK, Kim CW. Exosome derived from epigallocatechin gallate treated breast cancer cells suppresses tumor growth by inhibiting tumor-associated macrophage infiltration and M2 polarization. *BMC Cancer* 2013;13:421.
43. Yu P. Anti-inflammatory activitier of Praeruptorins and their molecular mechanisms of action [dissertation]. Guangdong: School of Pharmacy, Southern Medical University; 2013.
44. Fritz H, Kennedy DA, Ishii M, Fergusson D, Fernandes R, Cooley K, et al. Polysaccharide K and Coriolus versicolor extracts for lung cancer: a systematic review. *Integr Cancer Ther* 2015;14:201-211.
45. Shao B, Xu W, Dai H, Tu PF, Li HJ, Gao XM. A study on the immune receptors for polysaccharides from the roots of *Astragalus membranaceus*, a Chinese medicinal herb. *Biochem Biophys Res Commun* 2004;320:1103-1111.
46. Sekhon BK, Sze DM, Chan WK, Fan K, Li GQ, Moore DE, et al. PSP activates monocytes in resting human peripheral blood mononuclear cells: immunomodulatory implications for cancer

- treatment. Food Chem 2013;138:2201-2209.
47. Ghosh D, Maiti TK. Effects of native and heat-denatured Abrus agglutinin on tumor-associated macrophages in Dalton's lymphoma mice. Immunobiology 2007;212:667-673.
48. Woo SM, Choi YK, Cho SG, Sunju Park SJ, Ko SG. A new herbal formula, KSG-002, suppresses breast cancer growth and metastasis by targeting NF- κ B-dependent TNF α production in macrophages. Evid Based Complement Alternat Med 2013;1-10.
49. Qian X. Intervention of QHBJ prescription with the biological effects of COX-2/5-LOX metabolic pathway of arachidonic acid on ovarian carcinogenesis induced by DMBA in rats [dissertation]. Hangzhou: College of Pharmaceutical Sciences, Zhejiang University; 2012.
50. Lin X, Yi Z, Diao J, Shao M, Zhao L, Cai HB, et al. Shaoyao Decoction ameliorates colitis-associated colorectal cancer by downregulating proinflammatory cytokines and promoting epithelial-mesenchymal transition. J Transl Med 2014;12:105.
51. Lam W, Jiang Z, Guan J, Huang X, Hu R, Wang J, et al. PHY906 (KD018), an adjuvant based on a 1800-year-old Chinese medicine, enhanced the anti-tumor activity of Sorafenib by changing the tumor microenvironment. Sci Rep 2015;5:9384.
52. Liu Y, Yao F, Chai Y, Dong N, Sheng ZY, Yao YM. Xuebijing Injection promotes M2 polarization of macrophages and improves survival rate in septic mice. Evid Based Complement Alternat Med 2015;2015:352642.
53. Pollard JW. Tumour-educated macrophages promote tumour progression and metastasis. Nat Rev Cancer 2004;4:71-78.
54. Kurahara H, Shinichi H, Matakai Y, Maemura K, Noma H, Kubo F, et al. Significance of M2-polarized tumor-associated macrophage in pancreatic cancer. J Surg Res 2011;167:e211-e219.
55. Lewis CE, Pollard JW. Distinct role of macrophages in different tumor microenvironments. Cancer Res 2006;66:605-612.
56. Mantovani A, Romero P, Palucka AK, Marincola FM. Tumour immunity: effector response to tumour and role of the microenvironment. Lancet 2008;371:771-783.
57. Sica A, Mantovani A. Macrophage plasticity and polarization: *in vivo* veritas. J Clin Invest 2012;122:787-795.
58. Wang X. Research progress on the phenotype reversion of tumor-associated macrophage. Chin J Cancer Biother (Chin) 2014;21:216-217.
59. Iwanowycz S, Wang J, Jia X, Fan DP. Emodin inhibits breast cancer growth by modulating the phenotype of tumor-associated macrophages (TUM6P. 1004). J Immunol 2015;194:141-148.
60. Jia X, Yu F, Wang J, Iwanowycz S, Saaoud F, Wang Y, et al. Emodin suppresses pulmonary metastasis of breast cancer accompanied with decreased macrophage recruitment and M2 polarization in the lungs. Breast Cancer Res Treat 2014;148:291-302.
61. Zhao H, Zhang X, Chen X, Li Y, Ke ZQ, Tang T, et al. Isoliquiritigenin, a flavonoid from licorice, blocks M2 macrophage polarization in colitis-associated tumorigenesis through downregulating PGE₂ and IL-6. Toxicol Appl Pharmacol 2014;3:311-321.
62. Zhang H, Ren Y, Tang X, Wang K, Liu Y, Zhang L, et al. Vascular normalization induced by sinomenine hydrochloride results in suppressed mammary tumor growth and metastasis. Sci Rep 2015;5:8888.
63. Nguyen HYT, Vo BHT, Nguyen LTH, Bernad J, Alaeddine M, Coste A, et al. Extracts of *Crinum latifolium* inhibit the cell viability of mouse lymphoma cell line EL4 and induce activation of anti-tumour activity of macrophages *in vitro*. J Ethnopharmacol 2013;149:75-83.
64. Jiang Z, Chen Z, Li X, Zhao J, Li SM, Hu JP, et al. Immunomodulatory effects of *Sarcandra glabra* polysaccharides on macrophage RAW 264.7. Chin J Exp Tradit Med Form (Chin) 2014;20:178-182.
65. Jia C, Li F, He L, Li J. Effect of Fuzheng Jiedu Formula on tumor associated macrophage and related cytokine in the mouse gastric cancer postoperative recurrence model. Chin J Basic Med Tradit Chin Med (Chin) 2014;20:748-751.
66. Zhu R, Wu Y, Luo J, Yi L, Dong Y. Effect of berberine on mice RAW264.7 macrophages polarization. J Guangzhou Univ Tradit Chin Med (Chin) 2014;31:974-978.
67. Gao S, Zhou J, Liu N, Wang LJ, Gao QY, Wu Y, et al. Curcumin induces M2 macrophage polarization by secretion IL-4 and/or IL-13. J Mol Cell Cardiol 2015;85:131-139.
68. Hyam SR, Lee IA, Gu W, Kim KA, Jeong JJ, Jang SE, et al. Arctigenin ameliorates inflammation *in vitro* and *in vivo* by inhibiting the PI3K/AKT pathway and polarizing M1 macrophages to M2-like macrophages. Eur J Pharmacol 2013;708:21-29.
69. Solinas G, Germano G, Mantovani A, Allavena P. Tumor-associated macrophages (TAM) as major players of the cancer-related inflammation. J Leukocyte Biol 2009;86:1065-1073.
70. Mantovani A, Sozzani S, Locati M, Allavena P, Sica A. Macrophage polarization: tumor-associated macrophages as a paradigm for polarized M2 mononuclear phagocytes. Trends Immunol 2002;23:549-555.
71. Bingle L, Lewis CE, Corke KP, Reed MWR, Brown NJ. Macrophages promote angiogenesis in human breast tumour spheroids *in vivo*. Brit J Cancer 2006;94:101-107.
72. Egeblad M, Werb Z. New functions for the matrix metalloproteinases in cancer progression. Nat Rev Cancer 2002;2:161-174.
73. Steidl C, Lee T, Shah SP, Farinha P, Han G, Nayar T, et al. Tumor-associated macrophages and survival in classic Hodgkin's lymphoma. N Engl J Med. 2010; 362: 875-85.
74. Ito M, Ishii G, Nagai K, Maeda R, Nakano Y, Ochiai A. Prognostic impact of cancer-associated stromal cells in patients with stage I lung adenocarcinoma. CHEST J 2012;142:151-158.
75. Kurahara H, Takao S, Kuwahata T, Nagai T, Ding Q, Maeda K, et al. Clinical significance of folate receptor beta-expressing tumor-associated macrophages in pancreatic cancer. Ann Surg Oncol 2012;19:2264-2271.
76. Ding J, Xia Y, Liu C, Xu JY. Expression and clinical significance of M2 tumor-associated macrophage in pancreatic carcinoma. Cancer Res Prev Treat 2012;39:59-61.
77. Khan MA, Assiri AM, Broering DC. Complement and macrophage crosstalk during process of angiogenesis in tumor progression. J Biomed Sci 2015;22:58.
78. Riabov V, Gudima A, Wang N, Mickley A, Orekhov A, Kzyshkowska J. Role of tumor associated macrophages in tumor angiogenesis and lymphangiogenesis. Front Physiol 2014;5:75.
79. Kimura Y, Sumiyoshi M. Anti-tumor and anti-metastatic actions of wogonin isolated from *Scutellaria baicalensis* roots through anti-lymphangiogenesis. Phytomedicine 2013;20:328-336.
80. Zhang X. Study of depressant effect of PAMD to Expc-3 and the influence about the expression of tumorous new vessels in tumor-bearing athymic mouse [dissertation]. Harbin: Heilongjiang University of Chinese Medicine; 2009.
81. Varinska L, Gal P, Mojziso G, Mirossay L, Mojzis J. Soy and breast cancer: focus on angiogenesis. Int J Molecul Sci 2015;16:11728-11749.
82. Guo YP, Wang SH, Hoot DR, Clinton SK. Suppression of VEGF-mediated autocrine and paracrine interactions between prostate cancer cells and vascular endothelial cells by soy isoflavones. J Nutr Biochem 2007;18:408-417.
83. Ouyang H. Study on inhibitory effects and mechanism of QYHJ Decoction on human pancreatic cancer cell line CFPAC-1 in nude mice [dissertation]. Shanghai: Fudan University; 2009.
84. Yin J, Shi W, Zhu X, Chen Z, Liu LM. Qingyihuaji Formula inhibits progress of liver metastases from advanced pancreatic cancer xenograft by targeting to decrease expression of Cyr61 and VEGF. Integr Cancer Ther 2012;11:37-47.
85. Shi W. Establishment of human pancreatic carcinoma SW1990 cell

- Line with highly metastatic potential in the liver and interventional research of traditional Chinese drugs [dissertation]. Shanghai: Fudan University; 2007.
86. Lei Y, Guo X, Liu L, Zhou F. Research of *Rhodiola rosea* as the assistant anti-tumor drug. *Herald Med* 2014;33:1344-1347.
 87. Shi S, Wen P, Xu X. Anti-proliferation and anti-metastasis effects of emodin on human pancreatic cancer via NF- κ B signaling pathway. *Chin Arch Tradit Chin Med (Chin)* 2014;32:897-900.
 88. Lin Q. Bioactivity analysis of constituents from water caltrop pericarps and surveys about their anti-gastric cancer mechanism [dissertation]. Hangzhou: College of Biosystems Engineering and Food Science, Zhejiang University; 2013.
 89. Suboj P, Babykutty S, Valiyaparambil Gopi DR, Nair RS, Srinivas P, Gopala S. Aloe emodin inhibits colon cancer cell migration/angiogenesis by downregulating MMP-2/9, RhoB and VEGF via reduced DNA binding activity of NF-kappaB. *Eur J Pharm Sci* 2012;45:581-591.
 90. Deng J. Experimental study of the effect of antitumor and the molecular mechanism of honokiol [dissertation]. Hangzhou: Medical College of Zhejiang University; 2008.
 91. Schutyser E, Struyf S, Proost P, Opdenakker G, Laureys G, Verhasselt B, et al. Identification of biologically active chemokine isoforms from ascitic fluid and elevated levels of CCL18/pulmonary and activation-regulated chemokine in ovarian carcinoma. *J Biol Chem* 2002;277:24584-24593.
 92. Schepetkin IA, Quinn MT. Botanical polysaccharides: macrophage immunomodulation and therapeutic potential. *Intl Immunopharmacol* 2006;6:317-333.
 93. Liu C, Leung MY, Koon JC, Zhu LF, Hui YZ, Yu B, et al. Macrophage activation by polysaccharide biological response modifier isolated from *Aloe vera* L. var. *chinensis* (Haw.) Berg. *Int Immunopharmacol* 2006;6:1634-1641.
 94. Ye S, Zeng Y, Yin L. Effects of salidroside on proliferation, apoptosis, phagocytosis, ROS and NO production of murine peritoneal macrophages *in vitro*. *Chin J Cell Mol Immunol (Chin)* 2011;27:237-241.
 95. Li J, Xue X, Liu X, Fang J, Xie W. Effect of berberine on human endometrial carcinoma cell line Ishikawa co-cultured with TAM. *J Beijing Univ Tradit Chin Med (Chin)* 2011;34:448-452.
 96. Song Y, Ding N, Kanazawa T, Yamashita U, Yoshida Y. Cucurbitacin D is a new inflammasome activator in macrophages. *Int Immunopharmacol* 2013;17:1044-1050.
 97. Kim MY, Cho JY. 20S-dihydroprotoxanthadiol, a ginsenoside derivative, boosts innate immune responses of monocytes and macrophages. *J Ginseng Res* 2013;37:293-299.
 98. Dong Q, Sugiura T, Toyohira Y, Yoshida Y, Yangihara N, Karasaki Y. Stimulation of IFN-gamma production by garlic lectin in mouse spleen cells: involvement of IL-12 via activation of p38 MAPK and ERK in macrophages. *Phytomedicine* 2011;18:309-316.
 99. Shin JY, Song JY, Yun YS, Yang HO, Rhee DK, Pyo S. Immunostimulating effects of acidic polysaccharides extract of *Panax ginseng* on macrophage function. *Immunopharmacol Immunotoxicol* 2002;24:469-482.
 100. Diao H, Li X, Chen J, Luo Y, Chen X, Dong L, et al. *Bletilla striata* polysaccharide stimulates inducible nitric oxide synthase and proinflammatory cytokine expression in macrophages. *J Biosci Bioeng* 2008;105:85-89.
 101. Zhan X, Jia L, Niu Y, Qi HX, Chen XP, Zhan QW, et al. Targeted depletion of tumour-associated macrophages by an alendronate-glucosaminoglycan conjugate for cancer immunotherapy. *Biomaterials* 2014;35:10046-10057.
 102. Li X, Xu W. TLR4-mediated activation of macrophages by the polysaccharide fraction from *Polyporus umbellatus* (Pers.) Fries. *J Ethnopharmacol* 2011;135:1-6.
 103. Zhao Y. Traditional uses, phytochemistry, pharmacology, pharmacokinetics and quality control of *Polyporus umbellatus* (Pers.) Fries: a review. *J Ethnopharmacol* 2013;149:35-48.
 104. Li F, Huang D, Jiang L. Immunoregulatory effect of polysaccharide from the seeds of *Plantago asiatica* L. on RAW264.7 cells stimulated with lipopolysaccharide. *Food Sci* 2014;35:249-252.
 105. Zhu XL, Chen AF, Lin ZB. *Ganoderma lucidum* polysaccharides enhance the function of immunological effector cells in immunosuppressed mice. *J Ethnopharmacol* 2007;111:219-226.
 106. Lin Z. Antagonist effects of *Ganoderma polysaccharides* on tumor evasion from immune surveillance. *Chin J Pharmacol Toxicol (Chin)* 2015;29:14.
 107. Banerjee S, Parasramka M, Paruthy SB, eds. Polysaccharides in cancer prevention: from bench to bedside. Polysaccharides: Bioact Biotechnol 2015:2179-2214.
 108. Cai Z, Li W, Wang H, Yan WQ, Zhou YL, Wang GJ, et al. Anti-tumor and immunomodulating activities of a polysaccharide from the root of *Sanguisorba officinalis* L. *Int J Biol Macromolec* 2012;51:484-488.
 109. Tang X, Yan L, Gao J, Yang XL, Xu YX, Ge HY, et al. Antitumor and immunomodulatory activity of polysaccharides from the root of *Limonium sinense* Kuntze. *Int J Biol Macromolec* 2012;51:1134-1139.
 110. Sun X, Gao R, Xiong Y, Huang QC, Xu M. Antitumor and immunomodulatory effects of a water-soluble polysaccharide from *Lilii Bulbus* in mice. *Carbohydr Polym* 2014;102:543-549.
 111. Zhang J, Tang Q, Zhou C, Jia W, Da Siva L, Nguyen LD, et al. GLIS, a bioactive proteoglycan fraction from *Ganoderma lucidum*, displays anti-tumor activity by increasing both humoral and cellular immune response. *Life Sci* 2010;87:628-637.
 112. Nie X, Shi B, Ding Y, Tao W. Antitumor and immunomodulatory effects of Weikangfu Granule compound in tumor-bearing mice. *Curr Ther Res Clin Exp* 2006;67:138-150.
 113. Wang J, Tong X, Li P, Cao H, Su WW. Immuno-enhancement effects of Shenqi Fuzheng Injection on cyclophosphamide-induced immunosuppression in Balb/c mice. *J Ethnopharmacol* 2012;139:788-795.
 114. Shu G, Zhao W, Yue L, Su HW, Xiang MX. Antitumor immunostimulatory activity of polysaccharides from *Salvia chinensis* Benth. *J Ethnopharmacol* 2015;168:237-247.
 115. Kawanishi T, Ikeda-Dantsuji Y, Nagayama A. Effects of two basidiomycete species on interleukin 1 and interleukin 2 production by macrophage and T cell lines. *Immunobiology* 2010;215:516-520.
 116. Wang H, Chan Y, Li T, Wu C. Improving cachectic symptoms and immune strength of tumour-bearing mice in chemotherapy by a combination of *Scutellaria baicalensis* and Qing-Shu-Yi-Qi-Tang. *Eur J Cancer* 2012;48:1074-1084.
 117. Tu Y. The discovery of artemisinin (qinghaosu) and gifts from Chinese medicine. *Nat Med* 2011;17:1217-1220.
 118. Hingorani SR, Potter JD. Pancreas cancer meets the Thunder God. *Sci Transl Med* 2012;4:156ps21.
 119. Liu Z, Ma L, Zhou G-B. The main anticancer bullets of the Chinese medicinal herb, Thunder God vine. *Molecules* 2011;16:5283.
 120. Aggarwal BB, Gehlot P. Inflammation and cancer: how friendly is the relationship for cancer patients? *Curr Opin Pharmacol* 2009;9:351-369.
 121. Zhao FR, Mao HP, Zhang H, Hu LM, Wang H, Wang YF, et al. Antagonistic effects of two herbs in Zuojin Wan, a traditional Chinese medicine formula, on catecholamine secretion in bovine adrenal medullary cells. *Phytomedicine* 2010;17:659-668.
 122. Graziose R, Lila MA, Raskin I. Merging traditional Chinese medicine with modern drug discovery technologies to find novel drugs and functional foods. *Curr Drug Discov Technol* 2010;7:2-12.

(Received May 30, 2017)

Edited by YUAN Lin